

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/99775/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Liu, Zihao, Sanders, Andrew J ORCID: <https://orcid.org/0000-0002-7997-5286>,
Liang, Gehao, Song, Erwei, Jiang, Wen G ORCID: <https://orcid.org/0000-0002-3283-1111> and Gong, Chang 2017. Hey factors at the crossroad of
tumorigenesis and clinical therapeutic modulation of Hey for anticancer
treatment. *Molecular Cancer Therapeutics* 16 (5) , pp. 775-786. 10.1158/1535-7163.MCT-16-0576 file

Publishers page: <http://dx.doi.org/10.1158/1535-7163.MCT-16-0576>
<<http://dx.doi.org/10.1158/1535-7163.MCT-16-0576>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





Hey Factors at the Crossroad of Tumorigenesis and Clinical Therapeutic Modulation of Hey for Anticancer Treatment

Zihao Liu¹, Andrew J. Sanders², Gehao Liang¹, Erwei Song¹, Wen G. Jiang², and Chang Gong^{1,2}

Abstract

Hairy and Enhancer-of-split related with YRPW motif (Hey) transcription factors are important regulators of stem cell embryogenesis. Clinical relevance shows that they are also highly expressed in malignant carcinoma. Recent studies have highlighted functions for the Hey factors in tumor metastasis, the maintenance of cancer cell self-renewal, as well as proliferation and the promotion of tumor angiogenesis. Pathways which regulate Hey gene expression, such as Notch and TGFβ signaling, are frequently aberrant in numerous cancers. In addition, Hey factors control

downstream targets via recruitment of histone deacetylases (HDAC). Targeting these signaling pathways or HDACs may reverse tumor progression and provide clinical benefit for cancer patients. Thus, some small molecular inhibitors or monoclonal antibodies of each of these signaling pathways have been studied in clinical trials. This review focuses on the involvement of Hey proteins in malignant carcinoma progression and provides valuable therapeutic information for anticancer treatment. *Mol Cancer Ther*; 1–14. ©2017 AACR.

Introduction

Hairy and Enhancer-of-split related with YRPW motif factors (Hey1/2/L) belong to the basic helix-loop-helix Orange (bHLH-O) family which is also known as Hairy and Enhancer-of-split related protein (Hesr), Hairy-related transcriptional factor (HRT), Hes-related repressor protein (HERP), and cardiovascular helix-loop-helix factors (CHF; refs. 1–5). All three Hey genes have been found in developmental tissue, and abnormal expression of these proteins promotes abnormalities in stem cells, even leading to organ defects. Hey proteins can maintain an undifferentiated state of precursor cells by transcriptionally repressing cell fate regulators such as achaete-scute homolog 1 (6). In the developing heart, Hey proteins regulate cardiomyocyte precursor cell differentiation as well as epithelial-mesenchymal transition (EMT) of endocardium cells (7, 8). Since we recognized that cancer cells can monitor and utilize similar physiologic strategies to normal cells and promote tumor progression, for instance, cancer cells can initiate cellular plasticity and/or activate similar signaling pathways as mesenchymal cells, stem cells, or precursor cells do, we

have started to realize the significant role played by Hey factors in tumor progression (9, 10). Hey proteins are found to be selectively expressed in malignant tumor tissues, and numerous studies have been undertaken to explain the molecular mechanism governing the Hey proteins in tumorigenesis. The most outstanding feature is that many signaling pathways can potentially confer EMT via Hey factors in malignant carcinomas. In addition, Hey factors not only regulate differentiation, self-renewal, and proliferation of cancer cells, but contribute to tumor neovasculature as well. Accumulating evidence indicates Hey factors lay at the crossroad of tumor progression. However, there are currently very few review articles illustrating the roles of the Hey family in tumorigenesis. The current review explores the functional significance of the Hey family in initiating these processes. We also describe the signaling pathways involved in the control of Hey expression. Small molecular inhibitors or monoclonal antibodies to each of these signaling pathways show promising antitumor or antiangiogenic effect in clinical trials. Here, these promising avenues for cancer treatment are also discussed.

Structure of the Hey family proteins

Hey family members are highly conserved and resemble their homologs, the Hairy and Enhancer of Split (Hes) family, in the four domain structures: basic, helix-loop-helix (HLH), Orange, and two C-terminal motifs. Hey proteins are directly connected to the E-box DNA sequence (CANNTG) via the glycine-rich basic domain (11, 12). The bHLH-O domain serves as a platform for cofactor interaction (3, 13). Despite extensive homology with the Hes family, Hey proteins also have significant features that distinguish them from Hes proteins, namely, the YRPW motif (YHSW for HeyL) and GTEIGAF (GTEVGAF for Hey2) peptides (ref. 1; Fig. 1). Hey proteins have been regarded as transcription inhibitors in the past. They have since been shown to act as transcription activators as well as inhibitors (Table 1). Strikingly,

¹Guangdong Provincial Key Laboratory of Malignant Tumor Epigenetic and Gene Regulation, Breast Tumor Center, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, China. ²Cardiff China Medical Research Collaborative, Cardiff University School of Medicine, Cardiff University, Heath Park, Cardiff, United Kingdom.

Note: Supplementary data for this article are available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

Corresponding Authors: Chang Gong, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, 107 Yanjiang West Road, Guangzhou 510120, China. Phone: 8613925089353; Fax: 862081332853; E-mail: changgong282@163.com; or Wen G. Jiang, jiangW@cardiff.ac.uk

doi: 10.1158/1535-7163.MCT-16-0576

©2016 American Association for Cancer Research.




Figure 1
Domain structures with percentage identity of Hey2 and HeyL with Hey1. The bHLH domains show the highest conservation among other domains. The Orange domain shows less conservation. Individual Hey proteins potentially recruit selective cofactors via Orange domain and C-terminal motif.

their function seems to be regulated at multiple levels. For instance, nonsynonymous single-nucleotide polymorphism (SNP) naturally occurs at codon 94 of Hey1, which leads to a substitution of a leucine residue by methionine (L94M) in the helix 2 domain. The L94M-mutant Hey1 transforms from an androgen receptor corepressor to androgen receptor coactivator without changing its intrinsic repressive domains (14). The phosphorylation of the Serine-68 residue of Hey1 prevents its enhancement of p53 transcriptional activation but confers p53-activating chemotherapy resistance, whereas wild-type Hey1 stimulates p53 and alters the sensitivity to p53-activating chemotherapy drugs. Interestingly, such posttranscriptional regulation is also observed within Hey2 (15). The dynamic regulation of Hey proteins at pretranscriptional levels, posttranscriptional levels, or their own characteristic structure could partly explain why Hey proteins eliminate one target molecule in certain cancers but activate the same molecule or its analogues in others. It should be noted that the specificity for protein interactions and target molecules of different Hey variants is differential between certain cell types. L94M Hey1 variant strongly interacts with Hey2, whereas Hey1 forms an unstable homodimer with Hey2 (14). There is a potential, unknown Hey1 variant enhancing matrix metalloproteinase 9 (MMP9) expression in osteosarcoma, whereas wild-type Hey1 is unable to bind to the MMP9 promoter itself (16).

Hey proteins in malignant carcinomas

The levels of Hey factors are strikingly elevated in high-grade glioma, malignant osteosarcoma, high-grade esophageal squamous cell carcinoma, aggressive pancreatic adenocarcinoma rhabdomyosarcoma, as well as colorectal carcinoma (17–23). In these malignant carcinomas, aberrant Hey expression has been associated with poor prognosis, overall survival (OS), tumor grade, chemotherapy resistance, lymphatic metastasis, and vascular proliferative properties (17, 23–25). Taken together, these studies suggest that elevated levels of Hey proteins contribute to tumor progression, and to a certain extent, this is a result of their regulation of the behavior of cancer cells as well as remodeling of the tumor microenvironment (Fig. 2).


The roles of Hey proteins in cancer metastasis

It was first observed that Hey-induced EMT was required in the developing heart (26–29). Subsequently, Hey proteins were found to be involved in tumor metastasis progression. *In vitro*, Hey1 knockdown inhibited the invasive phenotype of osteosarcoma via downregulation of MMP9 (16). Furthermore, the transfection of Hey1 antisense oligonucleotides blocked EMT through E-cadherin expression, and Smad3 inhibition repressed the Hey1-induced EMT phenotype even with the presence of TGFβ (30). Strikingly, interaction between Hey1 and Smad3 has been observed *in vitro* (31), suggesting a Hey1–Smad3 complex transcriptionally represses E-cadherin. However, in the absence of activated Smad3, Hey1 does not influence EMT promotion, but only acts as a Snai1-initiated EMT marker (30, 32). On the other hand, Snai1, known as an E-cadherin repressor, potentially contributes to this repression process. Snai1 is reduced in Hey1/HeyL double knockouts and Hey2 knockout AV canals, and Snai1 can form a complex with Smad3 to occupy the E-cadherin promoter (26, 33). All these observations hint that Hey1 interacts with Smad3 and may inhibit E-cadherin directly or in a Snai1–Smad3–Hey1 manner. In other situations, Hey proteins promote mesenchymal–epithelial

Q8

Targets	Hey proteins	Comment	References
P53	Hey1, Hey2, HeyL	Activation	14, 15
MMP9	Hey1	Activation	16
Snai1	Hey1, Hey2, HeyL	Activation	26, 30
IL6	Hey1	Activation	37
Twist1	Hey1, Hey2	Repression	34
Snai2	Hey1, Hey2	Repression	34
Runx2	Hey1	Repression	40, 41
Col2a1	Hey1	Repression	42
VEGFR2	Hey1, Hey2	Repression	51, 58, 60, 61

Figure 2
Hey proteins in tumorigenesis. Via activating or inactivating cytokines and other transcriptional factors, Hey proteins show their regulation on tumor progression including cancer metastasis, cancer cells' quiescence maintenance as well as cancer neovasculation.



© 2017 American Association for Cancer Research

Molecular Cancer Therapeutics Reviews

AAGR

transition (MET). Upon Notch4 induction, Hey proteins promote melanoma MET and are important in promoting metastatic colonization because Hey1 and Hey2 can eliminate Snai2 as well as Twist1 expression via binding to their promoters (34). The different stimuli have a potential influence on Hey function, as TGFβ-induced Hey1 promotes EMT and Notch-induced Hey proteins regulate MET or transition irrelevant due to lack of Smad3. However, it is more complex than first thought. Forced expression of Hey proteins has no impact on Snai2 or E-cadherin expression in other cell lines (35, 36). Does the paradox happen in different cell types? Evidence from the previous section indicates the nonsynonymous SNP of Hey genes in different cell types will affect different Hey variants' DNA-binding ability as well as protein-interaction specificity. Based on this, we presume that Hey variants affect Snai1/Snai2 expression transcriptionally to mediate EMT/MET in different cell lines. More intensive research is required to fully characterize Hey variants and the posttranscriptional modification of Hey. Also, Hey1 participates in metastatic microenvironment remodeling. Tumor-derived Jagged1 enhances osteoblast secretion of IL6 via Hey1 activation, and, in turn, IL6 confers a

proliferative advantage to cancer cells (37). Epithelium-derived Jagged1 activates Hey1 which then promotes metastatic lymphoma cell chemotherapy resistance as well as progression in the tumor perivascular niche (38).

Hey proteins can regulate the differentiation, self-renewal, and proliferation of cancer cells

Hey proteins were identified as one of a few genes specifically expressed during embryogenesis (1, 39). Following this discovery, the potential capacity of the Hey family in sustaining cell quiescence was recognized (6, 40–42). Cancers monitor the quiescence strategy to keep their nondivide state and contribute to tumor progression (10, 43). The upregulation of Hey1 is likely to inhibit differentiation because rhabdomyosarcoma cells with an shRNA antagonizing Hey1 display differentiation morphology changes and the expression of differentiation marker myogenin (22). The introduction of Hey1 into proliferating osteosarcoma increases p53 expression and makes tumor cells stay in a nondividing state through p53-dependent reversible cell-cycle arrest (14). In the context of quiescence, elevated Hey family expression can reflect the undifferentiating property of malignant cancer cells. In

addition, the ability of Hey proteins in maintaining self-renewal was investigated. The expression of Hey1 and Hey2 is remarkably higher in cancer stem cells (CSC), also referred as tumor-initiating cells (TIC), than that in non-CSCs (44). Hey1 is supposed to maintain CSCs self-renewal capacity as the silencing of Hey1 dampens malignant tumor-initiating ability as well as tumor growth *in vivo* and reduces cancer cell sphere formation *in vitro* (45, 46). In hepatocellular carcinoma, Hey1 upregulation upon c-Met/FRA1 signaling increases the number and the size of TICs spheroids which represent the self-renewal ability of these cells (47). Furthermore, Hey proteins have an effect on cancer proliferation. Hey2 overexpression increases hepatocellular carcinoma cell viability and proliferation (48). HeyL can promote breast cancer initiation through interaction with TGF β -activated Smad3 (31). Interestingly, HeyL promotes p53-induced cell-cycle arrest which results in suppression of cancer cell proliferation and induction of cancer cell apoptosis in hepatocellular carcinoma (49). The same study also reported that 75% of hepatocellular carcinoma tissues had inactivation of HeyL, suggesting that HeyL is a potential tumor suppressor in hepatocellular carcinoma. This is an interesting observation, despite that it was a single study and demonstrated in a small cohort ($n = 80$), this will require further validation on a larger scale. However, the fact that HeyL differs in one of the key motifs, namely the YHSW motif, from Hey1 and Hey2 which both have the YRPW motif, may be one of the reasons why it acts differently from other Hey proteins. While YRPW appears to be a good target (16), YHSW, at least in hepatocellular carcinoma, may not be the case. This is clearly a fascinating area to explore, both in research and in clinical settings.

Balance between HeyL and Hey1/Hey2 regulates cancer neovasculature

Genetic studies have highlighted the great influence of Hey proteins in angiogenesis during development or pathologic conditions (27, 50–52). Angiogenesis actively requires a strict hierarchy between sprouting and vascular tubes (53). Previous research suggests a factor acting upstream of Hey, Delta-like 4 (DLL4), is capable of controlling this hierarchy, as the inhibition of DLL4 leads to a hyper-sprouting phenotype following exposure to proangiogenic factors (54–56). Much evidence, however, supports that DLL4's control on vascular sprouting is via its downstream factors, Hey1/2. It is acknowledged that epithelium with higher VEGFR2 emerges at the tip position and sustained VEGFR2 pathway activation results in excessive sprouting (57–59). Strikingly, Hey1, as well as Hey2, can suppress VEGFR2 expression and eliminate the increased frequency of epithelial cells at the tip position (58, 60, 61). When activated by the bone morphogenetic protein (BMP)/Activin receptor-like kinase (ALK) pathway, Hey1 as well as Hey2 abrogate the hyper-sprouting phenotype and induces tube formation (58, 62). In tumors, the coordinated balance between VEGFR2 and DLL4/Hey is tightly required for tumor cell expansion (63). DLL4/Hey2 overexpression leads to tumor growth by promoting low-density and mature tumor vessels through downregulation of VEGFR2 levels (64). DLL4/Hey blockage leads to VEGFR2 upregulation, which restrains tumor progression by producing hyper-sprouting, thin, fragile, and nonfunctional tumor vessels (56, 65–67). Interestingly, Jagged1-associated Hey upregulation seems to have little effect on low-density and mature tumor vessel phenotype, and Jagged1 promotes tumor-spouting angio-

genesis through distinct mechanisms (54, 68, 69). In contrast, HeyL potentially promotes neovascularization. Studies reveal that breast tumor-derived vascular samples exhibit at least 20-fold higher levels of HeyL than normal breast vasculature. The elevated level of HeyL potentially promotes neovasculture by forcing vascular endothelial cells to undergo proliferation and ceasing apoptosis (25). Taken together, this evidence highlights the complexity of Hey in angiogenesis, and drugs targeting DLL4, Jagged1, and ALK1 are promising.

Notch-Hey signaling pathway

The mature heterodimeric Notch receptors are cleaved at two sites once the five ligands (Delta-like 1, 3 and 4, and Jagged 1 and 2) bind to the four membrane-bound Notch receptors (Notch 1, 2, 3, and 4), firstly by a disintegrin and metalloprotease domain-containing protein 10/17 (ADAM10/17) and secondly by γ -secretase to release the Notch intracellular domain (NICD). In the nucleus, NICD interacts with the CBF1/Suppressor of Hairless/Lag1 (CSL) and recruits coactivators, allowing for transcriptional activation of Hey genes (4, 70, 71). Intriguingly, Notch receptors or Notch ligands show little selectivity for the induction of individual Hey proteins. Aberrant Notch-Hey axis shows great relevance to cancer biology. The Notch-Hey1 signaling pathway is over activated in invasive breast cell lines. Upon Notch inhibition via γ -secretase inhibitors (GSI), their migration and invasion capacity is reduced and this is accompanied by downregulation of Hey proteins (32, 72). The disruption of Notch-Hey1 in stroma bone cells decreases Jagged1-mediated breast tumor growth and bone metastasis (37). In osteosarcoma as well as rhabdomyosarcoma, Notch-Hey inhibition reverses tumor cell proliferative and relieves tumor burden (20, 22). GSI treatment also contributes to depletion of breast CSCs (44). Furthermore, a nonfunctional vascular network which results in tumor growth inhibition emerges when the DLL4-Notch-Hey1/2 pathway is blocked by DLL4 antibodies (56, 67). Thus, because GSIs, anti-Notch receptors, as well as anti-DLL4 are effective in Notch-Hey pathway inhibition, they have been developed into promising preclinical drugs (as summarized in Table 2).

γ -secretase inhibitors

Various preclinical trials show that GSIs have strong antitumor effects (73, 74). When treated with MK-0752 in phase I studies, clinical benefits such as complete response (CR) and prolonged stable disease (SD) were observed (75–78). However, patients present no objective responses to monotherapy of RO-4929097 in phase II clinical trials of solid tumors (79–82). Clinical indication of GSIs is still controversial, as a portion of cancer patients experienced SD during RO-4929097 or MK-0752 therapy, 1 advanced thyroid cancer patient achieved CR, and 71.4% (5/7) desmoid tumor patients had a partial response (PR) when they received another GSI, PF-0308414 (83). The most prominent and dose-limiting toxicity of GSIs is gastrointestinal (GI) events including diarrhea, vomiting, and nausea. This GI toxicity is likely based on the mechanism that inhibition of Notch signaling abrogates the undifferentiated state of intestinal crypt progenitor cells and results in differentiation into goblet cells (84). To reverse GI events, some investigators use glucocorticoid or tamoxifen therapy which potentially protects the intestine from goblet cell metaplasia (85, 86). Besides, the adverse events are scheduled dependent. Once-per-week dosing schedule of MK-0752 shows less severe GI events as well as fatigue than intermittent dosing for

Table 2. Selected therapeutic inhibitors of Notch signaling, TGF β signaling, and HDACs

Mechanism of action	Agent	Biology targeted	Clinical benefits	Disease	Stage	NCT number		
Notch								
Pan-Notch inhibitor	RO-4929097	Antitumor	SD, PD	Metastatic colorectal cancer	Phase II	NCT01116687		
			SD, PD	Recurrent ovarian cancer	Phase II	NCT01175343		
			SD	Pretreated pancreatic adenocarcinoma	Phase II	NCT01232829		
			PR, SD	Metastatic melanoma	Phase II	NCT01120275		
	MK-0752	Antitumor	CR, SD	Advanced solid tumors	Phase I	NCT00106145		
			SD	Children CNS malignancies	Phase I	NCT00572182		
			LY900009	Antitumor	SD	Advanced cancers	Phase I	NCT01158404
			PF-0308414	Antitumor	CR, PR, SD	Advanced solid tumors	Phase I	NCT00878189
	Notch1-specific antibody	OMP-52M51	Antitumor	PR, SD	Solid tumors	Phase I	NCT01778439	
		Notch2/3-specific antibody	OMP-59R5	Antitumor	SD, PD	Untreated metastatic pancreatic cancer	Phase I	NCT01647828
DLL4-specific antibody	REGN421		Angiogenesis targeting	PR, SD	Advanced solid tumors	Phase I	NCT00871559	
	OMP-21M18	Angiogenesis targeting	PR	Pretreated solid tumors	Phase I	NCT00744562		
TGFβ								
ALK1-specific antibody	ACE-041	Angiogenesis targeting	PR, SD	Advanced solid tumors	Phase I	NCT00996957		
	PF-03446962	Angiogenesis targeting	PR, SD	Pretreated advanced solid tumors	Phase I	NCT00557856		
			SD	Pretreated urothelial cancer	Phase II	NCT01620970		
			SD	Advanced solid tumors	Phase I	NCT01337050		
ALK4/5/7 antibody	LY2157299	Antitumor	CR, PR, SD	Advanced cancer and glioma	Phase I	Unavailable ^a		
			SD	Advanced solid tumors	Phase I	NCT01722825		
HDAC								
Pan-HDAC inhibitors	Vorinostat	Antitumor		Cutaneous T-cell lymphoma	Approved			
	Belinostat	Antitumor		Peripheral T-cell lymphoma	Approved			
	Panobinostat	Antitumor		Multiple myeloma	Approved			
	Romidepsin	Antitumor		Cutaneous T-cell lymphoma	Approved			

Abbreviation: CNS, central nervous system

^aReference 121.

3 to 7 days or continuous daily dosing group and once-per-week group also achieved substantial Notch signaling inhibition (75). With glucocorticoid therapy and intermittent schedule, cancer patients are more tolerant to higher GSIs exposure and may associate with better outcomes. However, it is worth considering that GSIs have an off-target effect as γ -secretase cleaves more than 90 substrates (87). Strikingly, two types of GSIs reduce Notch1 but not Notch4 activity, suggesting some GSIs are receptors specific (88). In addition, different GSIs enjoy quite inequivalent pharmacokinetics. LY900009 is cleared by oxidation and amide hydrolysis, and its renal clearance is low, while semagacestat, an analogue of LY900009, mostly depends on renal clearance (89, 90). RO4929097 is cleared by auto-induction of cytochrome P450 family 3 subfamily A polypeptide 4 (CYP3A4), indicating that combination RO4929097 therapy with antitumor agents metabolized by CYP3A4 might show limit clinical utility (91). Furthermore, intravenous GSIs are under development (ref. 92; chemical structures of the unproved GSIs are available in Supplementary Data: Supplementary Figs. S1–S4).

Anti-Notch receptor antibodies

As GSIs are pan-Notch inhibitors, several antibodies were launched to block Notch receptors more specifically by binding with their extracellular-negative regulator region or ligand-binding domain. Preclinical data show antitumor, antiangiogenesis effects and decreasing CSCs frequency following treatment with these receptor-specific antibodies (93–95). Based on the success of Notch-specific antibodies, OMP-52M51 (anti-Notch1) and OMP-59R5 (anti-Notch2/3) have been studied in clinical trials. In a phase I study in solid tumors, the best response to OMP-52M51 was 2 patients with adenoid cystic carcinomas: one achieved PR; the other had SD for 290 days

and SD was also observed in other tumors (96). Untreated metastatic pancreatic cancer patients only present SD, whereas 75% (6/8) of untreated extensive-stage small-cell lung cancer patients achieve PR to OMP-59R5 monotherapy (97, 98). Anti-Notch receptor antibodies are attractive, as they still function even in Notch receptors carrying mutations, and some of these tumors carrying mutations may be highly sensitive to these antibodies (93).

Anti-DLL4 monoclonal antibodies

Considering the great importance that DLL4 exerts on tumor vessel formation, targeting DLL4 was used to target tumor angiogenesis in preclinical studies, and several DLL4-blocking monoclonal antibodies have also been used to target Notch-mediated tumor angiogenesis in clinical trials (64, 99). SD and PR were noted in 41% of patients with advanced solid tumors when treated with REGN421 (Enoticumab), a DLL4 monoclonal antibody, in a phase I trial (100). OMP-21M18 (Demiczumab), another anti-DLL4 monoclonal antibody, showed antitumor effect, and 40% of patients with solid tumors responded with a reduction in tumor size (101). Although promising and well tolerated, severe adverse events, including hemangiomas, bleeding episodes, increased levels of troponin I, and ventricular dysfunction, were observed. In addition, targeting Jagged1 seems to exhibit an alternative therapeutic strategy which requires further clinical data (102, 103).

Agents in preclinical stage

Other agents blocking Notch signaling are also under development. Soluble decoys, which are Notch receptor extracellular domains or Notch ligands fused with or without human IgG, compete with endogenous ligands and inhibit Notch signaling activation. Notch1 decoys consisting of certain

EGF-like repeats can interrupt Jagged-class-induced Notch uniquely and show antiangiogenesis as well as antitumor effects with limited adverse events *in vivo* (68). Soluble DLL1 or Jagged1 decoys can also block Notch signaling successfully (104, 105). Also, cell-permeable peptides interact with NICD-CSL and form a transcriptionally repressive complex which halts leukemia cell proliferation (106, 107).

TGF β -Hey signaling pathway

Recent evidence has documented that TGF β signaling induces Hey protein in a Notch-independent manner or through canonical Notch. Upon TGF β 1 activation, initiation of Hey is conducted by Smad3/Smad4 complex binding to Hey promoters at Smad-binding element core repeat (SCR) positions, and Hey gene activation is still observed when canonical Notch is abrogated by GSI (30). BMP9 protein activates Smad1/5/8 and directly stimulates Hey expression via a noncanonical Notch signaling pathway when it binds to TGF β type I receptor-ALK1 receptor (58, 62, 108). On the other hand, activation of the Hey family can be enhanced by synergy between TGF- β /BMP and Notch signaling. Smads physically interact with Notch-dependent NICD and synergistically activate transcription of Hey1, Hey2, and HeyL, when Smads are activated by BMP-ALK5/6 or TGF β 1 treatment (30, 109, 110). As with the Notch pathway, TGF β signaling is often elevated in tumors and contributes to tumor progression. Subsequent studies have indicated a crucial role for TGF β in EMT initiation, and tumors break free from their neighboring tissue to undergo metastasis through TGF β -induced EMT (111). Smad3 is of significant importance for Hey1-induced EMT as Smad3 is an integral molecule for repressing E-cadherin. In addition to metastasis, TGF β pathway activation has been linked to tumor angiogenesis. Upon BMP9 treatment, the ALK1-Hey signaling pathway forces epithelial cells to remain in a stalk cell state, resulting in tube induction and mature vessel phenotype (58). If the ALK1-Hey signaling pathway is abrogated through addition of the ALK1 inhibitor, K02288, a hyper-sprouting phenotype is induced *in vitro* and angiogenesis is disrupted *in vivo* (112). Thus, TGF β receptor inhibitors, which are potentially antitumor as well as antiangiogenesis drugs, have been applied in preclinical trials (as summarized in Table 2).

ALK1 blockers

Several ALK1 inhibitors have been studied in clinical trials. ACE-041 (Dalantercept), another ALK1 blocker, was tested in squamous cell carcinoma, non-small cell lung cancer, and intestinal adenocarcinoma and displayed antitumor activity in phase II clinical trial (113). No responses or PR to PF-03446962, an antibody targeting ALK1, was observed in hepatocellular carcinoma, urothelial cancer, colorectal cancer, malignant pleural mesothelioma, and other solid tumors (114–116). Three patients with metastatic hepatocellular carcinoma, metastatic clear cell renal carcinoma, and KRAS-mutant non-small cell lung cancer showed PR to PF-03446962 in another phase I trial (117). SD was observed among these four studies. Although only a very small part of patients have PR to anti-ALK1, further research is required into anti-ALK1. PR and SD were observed in portions of patients who still had lesions and cancer progression following VEGFR tyrosine kinase inhibitor (TKI) treatment. The combination of VEGFR TKI and ACE-041

results in a promising antiangiogenesis effect with marked tumor vascular disruption in xenograft models (118).

ALK5 inhibitor

LY2157299 (galunisertib), a small molecular inhibitor targeting the TGF β receptor I, was originally developed as an ALK5 inhibitor and proved to complement ALK4/7 inhibitors (119). LY2157299 exerts an anti-invasive effect rather than antiproliferative effect on hepatocellular carcinoma cells via repression of Smad2 and Smad3 phosphorylation (120). A total of 24.3% of patients with glioma had either CR or PR to LY2157299, and 26.7% showed SD to LY2157299 in a phase I trial (121). Interestingly, 80% of low-grade glioma patients with isocitrate dehydrogenase mutation received clinical benefits in this study, when given LY2157299 treatment. In addition, LY2157299 is well tolerated and safe without adverse cardiac events. However, LY2157299 shows limited antitumor effects in pancreatic tumors (122).

Hey mediates histone deacetylases

The mechanisms through which Hey factors regulate their downstream effectors might also provide promising strategies for anticancer treatment. Hey factors are known to repress the expression of their target genes through recruitment of cofactors (123). Through Hey-mediated transcriptional repression, cancer cells maintain their undifferentiation state. Hey1 transcriptionally represses myogenin expression to sustain embryonal rhabdomyosarcoma cells in an immature state (22). Heterodimers between Hes1 and Hey factors potentially silence achaete-scute homolog 1, which results in the maintenance of an undifferentiated state of tumors (124–126). Histone deacetylases (HDAC) are potentially involved in the repressive effects of Hey factors, as treatment with trichostatin A, a pan class I and II HDAC inhibitor, partially abrogates the repressive effect of Hey factors (127–129). It has been suggested that Hey factors can use their bHLH domain to recruit the mSin3-NCoR-HDAC1 complex or associate with SIRT1, a member of NAD⁺-dependent HDACs, and induce transcriptional repression (11, 127). Further research indicates that Hey-HDACs complexes reduced target gene expression by downregulation of histone H3 lysine 27 acetylation (H3K27ac), which represents active transcription (130). Conversely, the inhibition of HDACs can lead to accumulation of acetylated histones and results in active transcription of target genes which are expected to cause tumor differentiation and induction of apoptosis (131, 132). Because the expression of HDACs is required for tumor cell survival and maintenance of an undifferentiated state, HDAC inhibitors might provide a new antitumor strategy. However, the application of HDAC inhibitors remains paradoxical and should be studied in different types of cancer. The silencing of HDAC1 and/or HDAC2 can give rise to hematologic malignancy initiation (133). Knockout of HDAC3 impairs genome stability as well as integrity and results in hepatocellular cancer (134).

HDAC inhibitors

Five HDAC inhibitors have been approved for T-cell lymphoma treatment, vorinostat (MK0683), belinostat (PXD-101), panobinostat (LBH-589), and romidepsin (FK-228), by the FDA, and chidamide (CS055/HBI-8000) approved in China (ref. 135; as summarized in Table 2). These highlight the impact of HDAC inhibitors as antitumor agents. A great number of HDAC

inhibitors are currently in testing in different phases of trials, either combined with other antitumor chemotherapeutics or as monotherapies. A phase II study indicates that entinostat (SNDX-275/MS-275), an inhibitor of HDAC 1 and 3, brings clinical benefits (PR, CR, and SD) to 24% of Hodgkin lymphoma patients, and the median progression-free survival (PFS) as well as OS of these patients was 5.5 months and 25.1 months, respectively (136). Entinostat also shows antitumor effect in several clinical trials (137, 138). In estrogen receptor–positive breast cancer, the combination of exemestane with entinostat improves median PFS to 4.3 months and median OS to 28.1 months, whereas median PFS and OS is 2.3 and 19.8 months, respectively, in the exemestane plus placebo group (139). Other HDAC inhibitors, such as ITF2357, CHR-3996, and JNJ-26481585, have been studied and show promising antitumor effect (140–142).

Combination of therapies

The combination of therapies targeting TGF β , HDACs, and Notch pathways requires thorough investigation regarding their cross-talk in specific cancer settings. For example, Notch and TGF β have synergetic carcinogenic effects in lung carcinoma, head and neck squamous, esophageal adenocarcinoma, renal cell carcinoma, thyroid carcinoma, and breast cancer (31, 143–146). Because both TGF β and Notch signaling can activate Hey, the simultaneous inhibition of both pathways might result in better outcomes than blockade of either individually. Interestingly, inhibition of both Notch and TGF β cannot increase the synergetic effects on inhibition of cancer cell migration, but additional blockage of Notch attenuates cancer cell proliferation in TGF β -treated cells (145). This highlights that combination therapies may affect more than one angle. Besides, combination of ALK1 inhibitors and GSI shows promise in targeting tumor angiogenesis, as inhibition of both signaling pathways further abolishes angiogenesis when compared with the inhibition of each alone (58). However, there is little clinical trial data about the combination of Notch and TGF β inhibitors, and further insightful studies are required. In another instance, targeting both Hey levels and Hey activity concomitantly might prove advantageous in cancer treatment. As an example, Hey proteins exert their influence on tumor cells by recruiting HDACs; when combining GSI and vorinostat, glioma and melanoma cells show a decreased viability (147).

Another strategy is to combine molecular-targeted drugs with classical chemotherapies. The combination of GSIs, HDAC inhibitors, or TGF β inhibitors with cytotoxic agents results in a more effective therapy since the inhibition of these pathways has been observed to enhance cancer cell lines sensitive to chemotherapy (148–150). Some clinical trials have also established the efficacy of combination therapies. For example, when combined GSIs with cytotoxic chemotherapy, clinical benefits, such as PR and prolonged SD, are observed in solid cancer patients (73, 151, 152). Encouraging antitumor activity is noticed in a Notch2/3-specific antibody study. Treatment OMP-59R5 with etoposide/cisplatin or nab-paclitaxel/gemcitabine shows 100% (3/3) PR in small cell lung cancer and 35% (9/26) PR and 35% (9/26) SD in untreated metastasis pancreatic cancer, respectively (97, 98). HDAC inhibitors in combination with classical chemotherapy also lead to a stronger antitumor effect. For instance, 64% thymoma and thymic carcinoma patients show objective response to belinostat in combination with cisplatin, doxorubicin and cyclophosphamide

and vorinostat combined with fludarabine, mitoxantrone, and dexamethasone results in a 77.8% overall response rate in relapsed or refractory mantle cell lymphoma (153, 154). However, the combination of HDAC inhibitors with chemotherapy may lead to unacceptable toxicity and on times is less efficient (155–157).

Perspective in selectively targeting Hey proteins and bHLH factors

Because different tumors tend to upregulate Hey proteins via distinct pathways, targeting Hey proteins directly may bring about a higher response rate than blocking these pathways individually. Besides, targeting Hey proteins themselves may result in fewer side effects because the target genes of Notch, TGF β , and HDAC signaling pathways will be unaffected. To target Hey, we have to understand the mechanism of action of Hey. There are two possible mechanisms of transcriptional regulation mediated by Hey proteins. The first mechanism is E-box–dependent transcriptional regulation. Hey proteins bind to E-box via basic domain and form functional complex with other cofactors via HLH domain. A domain located between amino acids 47 and 122 is necessary (11, 158). The second mechanism is E-box independent. Hey interacts with DNA-binding proteins via HLH-O domain and performs as a cofactor. The critical domains locate in amino acids 47 to 76 and 111 to 291, which stride over bHLH and Orange domains (61, 159). Based on these, some small molecular inhibitors to antagonize Hey–DNA interaction and Hey–cofactor interaction might be promising. Dimer inhibitors from natural compounds were reported to disrupt the Hey homolog Hes1 dimerization (160). It is still possible to isolate small-molecule inhibitors targeting Hey. In addition, mutagenesis of Hes1 amino acid sequence in the basic domain does not decrease its dimerization-forming ability, but abrogates its transcriptional function (161, 162). Thus, we may construct high structural compatible Hey-dominant–negative peptides which can form inert complexes with Hey and block the three critical functional domains of Hey to disrupt their protein–protein and DNA–protein interfaces. The most successful example is designing stabilized, cell-permeable peptides which bind with NICD–CSL complex and prevent mastermind-like-1 interfacing to antagonize leukemia proliferation (107).

Human bHLH transcription factors contain over 200 members and can be divided into five classes based on phylogenetic analysis (163). Hey transcriptional factors belong to clade B, and other transcriptional factors, such as Twist1-2 (clade A), MyoD (clade C), Max (clade D), Myc (clade E), and hypoxia-induced factor (HIF, clade E), are also bHLH factors. From the mechanistic inhibitory action of bHLH, the bHLH inhibitors can be summarized into the following groups: preventing dimerization, preventing DNA binding, and preventing bHLH factors expression (164). For example, some small-molecule inhibitors were isolated to specifically inhibit Myc–Max dimerization and block the binding of Myc–Max to DNA without affecting other structure-like bHLH factors dimerization (165, 166). By using Myc bHLH–Zip domain fragments, researchers also discovered local conformational changes and formation of hydrophobic cavities at the specific peptide sequences of the fragments upon binding with these small-molecule inhibitors (167). This makes it possible to design specific inhibitors simply through protein sequence analysis because the small-molecule binding sites have certain peptide

sequence criteria. Also, HIF dimer inhibitors as well as HIF DNA-binding inhibitors have been reported (168, 169). In addition, dominant negative peptides mimicking the HLH domain show a significant impact on E2A dimerization (170). Peptides of MyoD which have a high affinity for Id1 can interrupt MyoD-Id1 interaction and exhibit antitumor effects *in vitro* (171).

Conclusion

Hey proteins, a subfamily of mammalian bHLH-O transcriptional factors, have been highly investigated in several research studies since they have been found to be overexpressed in aggressive tumors. Previous work has focused on their transcriptional repressive roles in the maintenance of the undifferentiated state. More recently, studies reveal novel characteristics of Hey proteins in the regulation of cancer metastasis and their influence on angiogenesis. This article offers insight into the significant roles of Hey proteins in tumorigenesis. Alternatively, therapeutic agents able to reverse aberrant Notch, TGF β , and HDACs levels have been evaluated in clinical trials, but treatment-associated toxicities are also observed. Targeting Hey factors may represent an opportunity for higher response rates but fewer side effects than treatment with GSI, TGF β blockers, and HDAC inhibitors. Attention should be drawn to the Hey family in drug design, and studies must be carried out to analyze outcomes using Hey-specific inhibitors in the future.

References

- Leimeister C, Externbrink A, Klamt B, Gessler M. Hey genes: A novel subfamily of hairy-and Enhancer of split related genes specifically expressed during mouse embryogenesis. *Mech Dev* 1999;85:173–7.
- Kokubo H, Lun Y, Johnson RL. Identification and expression of a novel family of bHLH cDNAs related to Drosophila hairy and enhancer of split. *Biochem Biophys Res Commun* 1999;260:459–65.
- Nakagawa O, Nakagawa M, Richardson JA, Olson EN, Srivastava D. HRT1, HRT2, and HRT3: A new subclass of bHLH transcription factors marking specific cardiac, somitic, and pharyngeal arch segments. *Dev Biol* 1999; 216:72–84.
- Iso T, Sartorelli V, Chung G, Shichinohe T, Kedes L, Hamamori Y. HERP, a new primary target of notch regulated by ligand binding. *Mol Cell Biol* 2001;21:6071–9.
- Sakata Y, Kamei CN, Nakagami H, Bronson R, Liao JK, Chin MT. Ventricular septal defect and cardiomyopathy in mice lacking the transcription factor CHF1/Hey2. *Proc Natl Acad Sci* 2002;99:16197–202.
- Sakamoto M, Hirata H, Ohtsuka T, Bessho Y, Kageyama R. The basic helix-loop-helix genes Hes1/Hey1 and Hes2/Hey2 regulate maintenance of neural precursor cells in the brain. *J Biol Chem* 2003;278:44808–15.
- Fischer A, Leimeister C, Winkler C, Schumacher N, Klamt B, Elmasri H, et al. Hey bHLH factors in cardiovascular development. *Cold Spring Harb Symp Quant Biol* 2002;67:63–70.
- Fischer A, Gessler M. Hey genes in cardiovascular development. *Trends Cardiovasc Med* 2003;13:221–6.
- Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol* 2014;15:178–96.
- Sang L, Roberts JM, Collier HA. Hijacking HES1: How tumors co-opt the anti-differentiation strategies of quiescent cells. *Trends Mol Med* 2010; 16:17–26.
- Iso T, Sartorelli V, Poizat C, Iezzi S, Wu H-Y, Chung G, et al. HERP, a novel heterodimer partner of HES/E (spl) in Notch signaling. *Mol Cell Biol* 2001;21:6080–9.
- Heisig J, Weber D, Englberger E, Winkler A, Kneitz S, Sung W-K, et al. Target gene analysis by microarrays and chromatin immunoprecipitation identifies HEY proteins as highly redundant bHLH repressors. *PLoS Genet* 2012;8:e1002728.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The authors acknowledge Cancer Research Wales and Cardiff University China Medical Scholarship for supporting the study.

Grant Support

C. Gong was supported by grants from the Natural Science Foundation of China (8123060, 81442009, 81621004, 81272893, 81472466, and 81672594); National Science Foundation of Guangdong Province (2014A03036003, S2012030006287, 2014A030310378, 2015B050501004, and 2016A050502018); and Elite Young Scholars Program of Sun Yat-sen Memorial Hospital (Y201401). E. Song was supported by the Natural Science Foundation of China (81490750); Guangzhou Science Technology and Innovation Commission (201508020008 and 201508020249); and Guangdong Science and Technology Department (2015B050501004). Parts of this work were supported by grants to C. Gong obtained from Translational medicine public platform of Guangdong Province (4202037); Guangdong Department of Science & Technology Translational Medicine Center grant (2011A080300002); Grant KLB09001 from the Key Laboratory of Malignant Tumor Gene Regulation and Target Therapy of Guangdong Higher Education Institutes, Sun Yat-Sen University; and Grant (2013) 163 from Key Laboratory of Malignant Tumor Molecular Mechanism and Translational Medicine of Guangzhou Bureau of Science and Information Technology.

Received October 2, 2016; revised December 29, 2016; accepted December 29, 2016; published OnlineFirst xx xx, xxxx.

- Davis RL, Turner DL. Vertebrate hairy and Enhancer of split related proteins: Transcriptional repressors regulating cellular differentiation and embryonic patterning. *Oncogene* 2001;20:8342–57.
- Villaronga MA, Lavery DN, Bevan CL, Llanos S, Beldandia B. HEY1 Leu94-Met gene polymorphism dramatically modifies its biological functions. *Oncogene* 2010;29:411–20.
- López-Mateo I, Arruabarrena-Aristorena A, Artaza-Irigaray C, López JA, Calvo E, Beldandia B. HEY1 functions are regulated by its phosphorylation at Ser-68. *Biosci Rep* 2016;36:e00343.
- Tsuru A, Setoguchi T, Matsunoshita Y, Nagao-Kitamoto H, Nagano S, Yokouchi M, et al. Hairy/enhancer-of-split related with YRPW motif protein 1 promotes osteosarcoma metastasis via matrix metalloproteinase 9 expression. *Br J Cancer* 2015;112:1232–40.
- Forghanifard MM, Taleb S, Abbaszadegan MR. Notch signaling target genes are directly correlated to esophageal squamous cell carcinoma tumorigenesis. *Pathol Oncol Res* 2015;21:463–7.
- Gaetani P, Hulleman E, Levi D, Quarto M, Scorsetti M, Helin K, et al. Expression of the transcription factor HEY1 in glioblastoma: A preliminary clinical study. *Tumori* 2010;96:97.
- El Hindy N, Keyvani K, Pagenstecher A, Dammann P, Sandalcioğlu IE, Sure U, et al. Implications of Dll4-Notch signaling activation in primary glioblastoma multiforme. *Neurooncology* 2013;15:1366–78.
- Engin F, Bertin T, Ma O, Jiang MM, Wang L, Sutton RE, et al. Notch signaling contributes to the pathogenesis of human osteosarcomas. *Hum Mol Genet* 2009;18:1464–70.
- Mullendore ME, Koorstra J-B, Li Y-M, Offerhaus GJ, Fan X, Henderson CM, et al. Ligand-dependent Notch signaling is involved in tumor initiation and tumor maintenance in pancreatic cancer. *Clin Cancer Res* 2009;15: 2291–301.
- Belyea BC, Naini S, Bentley RC, Linardic CM. Inhibition of the Notch-Hey1 axis blocks embryonal rhabdomyosarcoma tumorigenesis. *Clin Cancer Res* 2011;17:7324–36.
- Candy PA, Phillips MR, Redfern AD, Colley SM, Davidson JA, Stuart LM, et al. Notch-induced transcription factors are predictive of survival and 5-fluorouracil response in colorectal cancer patients. *Br J Cancer* 2013;109: 1023–30.

- 24 Hulleman E, Quarto M, Vernell R, Masserdotti G, Colli E, Kros JM, et al. A role for the transcription factor HEY1 in glioblastoma. *J Cell Mol Med* 2009;13:136–46.
- 25 Parker BS, Argani P, Cook BP, Liangfeng H, Chartrand SD, Zhang M, et al. Alterations in vascular gene expression in invasive breast carcinoma. *Cancer Res* 2004;64:7857–66.
- 26 Fischer A, Steidl C, Wagner TU, Lang E, Jakob PM, Friedl P, et al. Combined loss of Hey1 and HeyL causes congenital heart defects because of impaired epithelial to mesenchymal transition. *Circ Res* 2007;100:856–63.
- 27 Kokubo H, Miyagawa-Tomita S, Nakazawa M, Saga Y, Johnson RL. Mouse hesr1 and hesr2 genes are redundantly required to mediate Notch signaling in the developing cardiovascular system. *Dev Biol* 2005;278:301–9.
- 28 Rutenberg JB, Fischer A, Jia H, Gessler M, Zhong TP, Mercola M. Developmental patterning of the cardiac atrioventricular canal by Notch and Hairy-related transcription factors. *Development* 2006;133:4381–90.
- 29 Luna-Zurita L, Prados B, Grego-Bessa J, Luxan G, del Monte G, Benguria A, et al. Integration of a Notch-dependent mesenchymal gene program and Bmp2-driven cell invasiveness regulates murine cardiac valve formation. *J Clin Invest* 2010;120:3493–507.
- 30 Zavadil J, Cermak L, SotoNieves N, Böttinger EP. Integration of TGFβ/Smad and Jagged1/Notch signalling in epithelial to mesenchymal transition. *EMBO J* 2004;23:1155–65.
- 31 Han L, Diehl A, Nguyen NK, Korangath P, Teo W, Cho S, et al. The Notch pathway inhibits TGFβ signaling in breast cancer through HEYL-mediated crosstalk. *Cancer Res* 2014;74:6509–18.
- 32 Bolos V, Mira E, Martinez-Poveda B, Luxan G, Canamero M, Martinez AC, et al. Notch activation stimulates migration of breast cancer cells and promotes tumor growth. *Breast Cancer Res* 2013;15:R54.
- 33 Vincent T, Neve EPA, Johnson JR, Kukalev A, Rojo F, Albanell J, et al. A SNAIL1-SMAD3/4 transcriptional repressor complex promotes TGF-β mediated epithelial-mesenchymal transition. *Nat Cell Biol* 2009;11:943–50.
- 34 Bonyadi Rad E, Hammerlindl H, Wels C, Popper U, Ravindran Menon D, Breiteneder H, et al. Notch4 signaling induces a mesenchymal-epithelial-like transition in melanoma cells to suppress malignant behaviors. *Cancer Res* 2016;76:1690–7.
- 35 Leong KG, Niessen K, Kulic I, Raouf A, Eaves C, Pollet I, et al. Jagged1-mediated Notch activation induces epithelial-to-mesenchymal transition through Slug-induced repression of E-cadherin. *J Exp Med* 2007;204:2935–48.
- 36 Niessen K, Fu Y, Chang L, Hoodless PA, Mcfadden D, Karsan A. Slug is a direct Notch target required for initiation of cardiac cushion cellularization. *J Cell Biol* 2008;182:315–25.
- 37 Sethi N, Dai X, Winter CG, Kang Y. Tumor-derived JAGGED1 promotes osteolytic bone metastasis of breast cancer by engaging notch signaling in bone cells. *Cancer Cell* 2011;19:192–205.
- 38 Cao Z, Ding BS, Guo P, Lee SB, Butler JM, Casey SC, et al. Angiocrine factors deployed by tumor vascular niche induce B cell lymphoma invasiveness and chemoresistance. *Cancer Cell* 2014;25:350–65.
- 39 Leimeister C, Schumacher N, Steidl C, Gessler M. Analysis of HeyL expression in wild-type and Notch pathway mutant mouse embryos. *Mech Dev* 2000;98:175–8.
- 40 Zamurovic N, Cappellen D, Rohner D, Susa M. Coordinated activation of notch, Wnt, and transforming growth factor-beta signaling pathways in bone morphogenic protein 2-induced osteogenesis. Notch target gene Hey1 inhibits mineralization and Runx2 transcriptional activity. *J Biol Chem* 2004;279:37704–15.
- 41 Hilton MJ, Tu X, Wu X, Bai S, Zhao H, Kobayashi T, et al. Notch signaling maintains bone marrow mesenchymal progenitors by suppressing osteoblast differentiation. *Nat Med* 2008;14:306–14.
- 42 Grogan SP, Olee T, Hiraoka K, Lotz MK. Repression of chondrogenesis through binding of notch signaling proteins HES-1 and HEY-1 to N-box domains in the COL2A1 enhancer site. *Arthritis Rheum* 2008;58:2754–63.
- 43 Yeh AC, Ramaswamy S. Mechanisms of cancer cell dormancy—another hallmark of cancer? *Cancer Res* 2015;75:5014–22.
- 44 Yamamoto M, Taguchi Y, Ito-Kureha T, Semba K, Yamaguchi N, Inoue J. NF-kappaB non-cell-autonomously regulates cancer stem cell populations in the basal-like breast cancer subtype. *Nat Commun* 2013;4:2299.
- 45 Zhu P, Wang Y, Du Y, He L, Huang G, Zhang G, et al. C8orf4 negatively regulates self-renewal of liver cancer stem cells via suppression of NOTCH2 signalling. *Nat Commun* 2015;6:7122.
- 46 Wu H-C, Lin Y-C, Liu C-H, Chung H-C, Wang Y-T, Lin Y-W, et al. USP11 regulates PML stability to control Notch-induced malignancy in brain tumours. *Nat Commun* 2014;5.
- 47 Lau EY, Lo J, Cheng BY, Ma MK, Lee JM, Ng JK, et al. Cancer-associated fibroblasts regulate tumor-initiating cell plasticity in hepatocellular carcinoma through c-Met/FRA1/HEY1 signaling. *Cell Rep* 2016;15:1175–89.
- 48 Wu DC, Zhang MF, Su SG, Fang HY, Wang XH, He D, et al. HEY2, a target of miR-137, indicates poor outcomes and promotes cell proliferation and migration in hepatocellular carcinoma. *Oncotarget* 2016;7:38052–63.
- 49 Kuo KK, Jian SF, Li YJ, Wan SW, Weng CC, Fang K, et al. Epigenetic inactivation of transforming growth factorβ1 target gene HEYL, a novel tumor suppressor, is involved in the P53-induced apoptotic pathway in hepatocellular carcinoma. *Hepatol Res* 2015;45:782–93.
- 50 Fischer A, Schumacher N, Maier M, Sendtner M, Gessler M. The Notch target genes Hey1 and Hey2 are required for embryonic vascular development. *Genes Dev* 2004;18:901–11.
- 51 Li B, Tang SB, Hu J, Gao Y, Zhang G, Lin SF, et al. Protective effects of transcription factor HESR1 on retinal vasculature. *Microvasc Res* 2006;72:146–52.
- 52 Adepoju O, Wong A, Kitajewski A, Tong K, Boscolo E, Bischoff J, et al. Expression of HES and HEY genes in infantile hemangiomas. *Vasc Cell* 2011;3:19.
- 53 Tung JJ, Tattersall IW, Kitajewski J. Tips, stalks, tubes: Notch-mediated cell fate determination and mechanisms of tubulogenesis during angiogenesis. *Cold Spring Harbor Perspect Med* 2012;2:a006601.
- 54 Benedito R, Roca C, Sörensen I, Adams S, Gossler A, Fruttiger M, et al. The notch ligands Dll4 and Jagged1 have opposing effects on angiogenesis. *Cell* 2009;137:1124–35.
- 55 Liu Z, Fan F, Wang A, Zheng S, Lu Y. Dll4-Notch signaling in regulation of tumor angiogenesis. *J Cancer Res Clin Oncol* 2014;140:525–36.
- 56 Noguera-Troise I, Daly C, Papadopoulos NJ, Coetsee S, Bolland P, Gale NW, et al. Blockade of Dll4 inhibits tumour growth by promoting non-productive angiogenesis. *Nature* 2006;444:1032–7.
- 57 Jakobsson L, Franco CA, Bentley K, Collins RT, Ponsioen B, Aspö IM, et al. Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting. *Nat Cell Biol* 2010;12:943–53.
- 58 Larrivee B, Prahst C, Gordon E, del Toro R, Mathivet T, Duarte A, et al. ALK1 signaling inhibits angiogenesis by cooperating with the Notch pathway. *Dev Cell* 2012;22:489–500.
- 59 Boareto M, Jolly MK, Ben-Jacob E, Onuchic JN. Jagged mediates differences in normal and tumor angiogenesis by affecting tip-stalk fate decision. *Proc Natl Acad Sci* 2015;112:E3836–E44.
- 60 Taylor KL, Henderson AM, Hughes CCW. Notch activation during endothelial cell network formation in vitro targets the basic HLH transcription factor HESR-1 and downregulates VEGFR-2/KDR expression. *Microvasc Res* 2002;64:372–83.
- 61 Holderfield MT, Henderson Anderson AM, Kokubo H, Chin MT, Johnson RL, Hughes CC. HESR1/CHF2 suppresses VEGFR2 transcription independent of binding to E-boxes. *Biochem Biophys Res Commun* 2006;346:637–48.
- 62 Ricard N, Ciais D, Levot S, Subileau M, Mallet C, Zimmers TA, et al. BMP9 and BMP10 are critical for postnatal retinal vascular remodeling. *Blood* 2012;119:6162–71.
- 63 Siemerink MJ, Klaassen I, Van Noorden CJ, Schlingemann RO. Endothelial tip cells in ocular angiogenesis: Potential target for anti-angiogenesis therapy. *J Histochem Cytochem* 2013;61:101–15.
- 64 Li JL, Sainson RC, Shi W, Leek R, Harrington LS, Preusser M, et al. Delta-like 4 Notch ligand regulates tumor angiogenesis, improves tumor vascular function, and promotes tumor growth in vivo. *Cancer Res* 2007;67:11244–53.
- 65 Patel NS, Li JL, Generali D, Poulson R, Cranston DW, Harris AL. Up-regulation of delta-like 4 ligand in human tumor vasculature and the role of basal expression in endothelial cell function. *Cancer Res* 2005;65:8690–7.
- 66 Ridgway J, Zhang G, Wu Y, Stawicki S, Liang W-C, Chantry Y, et al. Inhibition of Dll4 signalling inhibits tumour growth by deregulating angiogenesis. *Nature* 2006;444:1083–7.

- 67 Djokovic D, Trindade A, Gigante J, Pinho M, Harris AL, Duarte A. Incomplete DLL4/Notch signaling inhibition promotes functional angiogenesis supporting the growth of skin papillomas. *BMC Cancer* 2015; 15:608.
- 68 Kangsamaksin T, Murtomaki A, Kofler NM, Cuervo H, Chaudhri RA, Tattersall IW, et al. NOTCH decoys that selectively block DLL/NOTCH or JAG/NOTCH disrupt angiogenesis by unique mechanisms to inhibit tumor growth. *Cancer Discov* 2015;5:182–97.
- 69 Qiu XX, Chen L, Wang CH, Lin ZX, Chen BJ, You N, et al. The vascular Notch ligands delta-like ligand 4 (DLL4) and Jagged1 (JAG1) have opposing correlations with microvascularization but uniform prognostic effect in primary glioblastoma: A preliminary study. *World Neurosurg* 2015;88:447–58.
- 70 Nakagawa O, McFadden DG, Nakagawa M, Yanagisawa H, Hu T, Srivastava D, et al. Members of the HRT family of basic helix–loop–helix proteins act as transcriptional repressors downstream of Notch signaling. *Proc Natl Acad Sci* 2000;97:13655–60.
- 71 Vinson KE, George DC, Fender AW, Bertrand FE, Sigounas G. The Notch pathway in colorectal cancer. *Int J Cancer* 2015;138:1835–42.
- 72 Chen J, Imanaka N, Griffin J. Hypoxia potentiates Notch signaling in breast cancer leading to decreased E-cadherin expression and increased cell migration and invasion. *Br J Cancer* 2010;102:351–60.
- 73 Schott AF, Landis MD, Dontu G, Griffith KA, Layman RM, Krop I, et al. Preclinical and clinical studies of gamma secretase inhibitors with docetaxel on human breast tumors. *Clin Cancer Res* 2013;19:1512–24.
- 74 Luistro L, He W, Smith M, Packman K, Vilenchik M, Carvajal D, et al. Preclinical profile of a potent gamma-secretase inhibitor targeting notch signaling with in vivo efficacy and pharmacodynamic properties. *Cancer Res* 2009;69:7672–80.
- 75 Krop I, Demuth T, Guthrie T, Wen PY, Mason WP, Chinnaiyan P, et al. Phase I pharmacologic and pharmacodynamic study of the gamma secretase (Notch) inhibitor MK-0752 in adult patients with advanced solid tumors. *J Clin Oncol* 2012;30:2307–13.
- 76 Fouladi M, Stewart CF, Olson J, Wagner LM, Onar-Thomas A, Kocak M, et al. Phase I trial of MK-0752 in children with refractory CNS malignancies: A pediatric brain tumor consortium study. *J Clin Oncol* 2011;29:3529–34.
- 77 Piha-Paul SA, Munster PN, Hollebecque A, Argilés G, Dajani O, Cheng JD, et al. Results of a phase I trial combining ridaforolimus and MK-0752 in patients with advanced solid tumours. *Eur J Cancer* 2015;51:1865–73.
- 78 Hoffman LM, Fouladi M, Olson J, Daryani VM, Stewart CF, Wetmore C, et al. Phase I trial of weekly MK-0752 in children with refractory central nervous system malignancies: a pediatric brain tumor consortium study. *Childs Nerv Syst* 2015;31:1283–9.
- 79 Diaz-Padilla I, Wilson MK, Clarke BA, Hirte HW, Welch SA, Mackay HJ, et al. A phase II study of single-agent RO4929097, a gamma-secretase inhibitor of Notch signaling, in patients with recurrent platinum-resistant epithelial ovarian cancer: A study of the Princess Margaret, Chicago and California phase II consortia. *Gynecol Oncol* 2015;137:216–22.
- 80 Jesus-Acosta AD, Laheru D, Maitra A, Arcaroli J, Rudek MA, Dasari A, et al. A phase II study of the gamma secretase inhibitor RO4929097 in patients with previously treated metastatic pancreatic adenocarcinoma. *Invest New Drugs* 2014;32:739–45.
- 81 Lee SM, James Moon MS, Do BGR, Tarek Chidiac MD, Flaherty LE, Zha Y, et al. Phase 2 study of RO4929097, a gamma-secretase inhibitor, in metastatic melanoma: SWOG 0933. *Cancer* 2015;121:432–40.
- 82 Strosberg JR, Yeatman T, Weber J, Coppola D, Schell MJ, Gang H, et al. A phase II study of RO4929097 in metastatic colorectal cancer. *Eur J Cancer* 2012;48:997–1003.
- 83 Messersmith WA, Shapiro GI, Cleary JM, Jimeno A, Dasari A, Huang B, et al. A phase I, dose-finding study in patients with advanced solid malignancies of the oral gamma-secretase inhibitor PF-03084014. *Clin Cancer Res* 2015;21:60–7.
- 84 van Es JH, van Gijn ME, Riccio O, van den Born M, Vooijs M, Begthel H, et al. Notch/ γ -secretase inhibition turns proliferative cells in intestinal crypts and adenomas into goblet cells. *Nature* 2005;435:959–63.
- 85 Samon JB, Castillo-Martin M, Hadler M, Ambesi-Impioabato A, Paietta E, Racevskis J, et al. Preclinical analysis of the γ -secretase inhibitor PF-03084014 in combination with glucocorticoids in T-cell acute lymphoblastic leukemia. *Mol Cancer Ther* 2012;11:1565–75.
- 86 Backus K. Crosstalk between PKC α and Notch-4 in endocrine-resistant breast cancer cells. *Oncogenesis* 2013;2:e60.
- 87 Langosch D, Steiner H. Substrate processing in intramembrane proteolysis by γ -secretase – the role of protein dynamics. *Biol Chem* 2016Dec 28. [Epub ahead of print].
- 88 Harrison H, Farnie G, Howell SJ, Rock RE, Stylianou S, Brennan KR, et al. Regulation of breast cancer stem cell activity by signaling through the Notch4 receptor. *Cancer Res* 2010;70:709–18.
- 89 Pant S, Jones SF, Kurkjian CD, Infante JR, Moore KN, Burris HA, et al. A first-in-human phase I study of the oral Notch inhibitor, LY900009, in patients with advanced cancer. *Eur J Cancer* 2016;56:1–9.
- 90 Yi P, Hadden C, Kulanthaivel P, Calvert N, Annes W, Brown T, et al. Disposition and metabolism of semagacestat, a γ -secretase inhibitor, in humans. *Drug Metab Dispos* 2010;38:554–65.
- 91 Tolcher AW, Messersmith WA, Mikulski SM, Papadopoulos KP, Kwak EL, Gibbon DG, et al. Phase I study of RO4929097, a gamma secretase inhibitor of Notch signaling, in patients with refractory metastatic or locally advanced solid tumors. *J Clin Oncol* 2012;30:2348–53.
- 92 Knoechel B, Bhatt A, Pan L, Pedamallu CS, Severson E, Gutierrez A, et al. Complete hematologic response of early T-cell progenitor acute lymphoblastic leukemia to the γ -secretase inhibitor BMS-906024: genetic and epigenetic findings in an outlier case. *Cold Spring Harbor Mol Case Studies* 2015;1:a000539.
- 93 Wallace B, Wang M, Muriel C, Cain J, Cancilla B, Shah J, et al. Novel NOTCH3 activating mutations identified in tumors sensitive to OMP-59R5, a monoclonal antibody targeting the Notch2 and Notch3 receptors [abstract]. In: Proceedings of the 104th Annual Meeting of the American Association for Cancer Research; 2013 Apr 6–10; Washington, DC. Philadelphia (PA): AACR. Abstract nr 213.
- 94 Asteamézaga M, Zhang N, Lineberger JE, Arnold BA, Toner TJ, Gu M, et al. Characterization of Notch1 antibodies that inhibit signaling of both normal and mutated Notch1 receptors. *PLoS One* 2010;5:e9094.
- 95 Yen WC, Fischer MM, Axelrod F, Bond C, Cain J, Cancilla B, et al. Targeting Notch signaling with a Notch2/Notch3 antagonist (tarextumab) inhibits tumor growth and decreases tumor-initiating cell frequency. *Clin Cancer Res* 2015;21:2084–95.
- 96 Munster P, Eckhardt SG, Patnaik A, Shields AF, Tolcher AW, Davis SL, et al. Safety and preliminary efficacy results of a first-in-human phase I study of the novel cancer stem cell (CSC) targeting antibody brontic-tuzumab (OMP-52M51, anti-Notch1) administered intravenously to patients with certain advanced solid tumors. *Mol Cancer Ther* 2015; 14 (12 Suppl 2):C42.
- 97 Bendell J, Cohn A, Smith L, Strickler J, Gluck W, Schmidt W, et al. 688P Final results of a phase 1B of OMP-59R5 (anti-notch2/3/stem cell antibody) in combination with nab-paclitaxel and gemcitabine (NAB-P+GEM) in patients (PTS) with untreated metastatic pancreatic cancer (MPC): Alpine study. *Ann Oncol* 2014;25:iv233–iv4.
- 98 Pietanza M, Spira A, Jotte R, Gadgeel S, Mita A, Liu S, et al. 1473P Phase 1B trial of anti-notch 2/3 antibody OMP-59R5 in combination with etoposide and cisplatin (EP) in patients (PTS) with untreated extensive-stage small-cell lung cancer (ED-SCLC): the pinnacle study. *Ann Oncol* 2014;25: iv514–iv5.
- 99 Scehnet JS, Jiang W, Kumar SR, Krasnoperov V, Trindade A, Benedito R, et al. Inhibition of DLL4-mediated signaling induces proliferation of immature vessels and results in poor tissue perfusion. *Blood* 2007;109: 4753–60.
- 100 Chiorean EG, Lorusso P, Strother RM, Diamond JR, Younger A, Messersmith WA, et al. A Phase I first-in-human study of enoticumab (REGN421), a fully human delta-like ligand 4 (DLL4) monoclonal antibody in patients with advanced solid tumors. *Clin Cancer Res* 2015;21: 2695–703.
- 101 Smith DC, Eisenberg PD, Manikhas G, Chugh R, Gubens MA, Staggs RJ, et al. A phase I dose escalation and expansion study of the anticancer stem cell agent demcizumab (anti-DLL4) in patients with previously treated solid tumors. *Clin Cancer Res* 2014;20:6295–303.
- 102 Steg AD, Katre AA, Goodman B, Han HD, Nick AM, Stone RL, et al. Targeting the notch ligand JAGGED1 in both tumor cells and stroma in ovarian cancer. *Clin Cancer Res* 2011;17:5674–85.
- 103 Chen JY, Li CF, Chu PY, Lai YS, Chen CH, Jiang SS, et al. Lysine demethylase 2A promotes stemness and angiogenesis of breast cancer by upregulating Jagged1. *Oncotarget* 2014;7:27689–710.

- 104 Varnum-Finney B, Wu L, Yu M, Brashem-Stein C, Staats S, Flowers D, et al. Immobilization of Notch ligand, Delta-1, is required for induction of notch signaling. *J Cell Sci* 2000;113:228–41.
- 105 Small D, Kovalenko D, Kacer D, Liaw L, Landriscina M, Di SC, et al. Soluble Jagged 1 represses the function of its transmembrane form to induce the formation of the Src-dependent chord-like phenotype. *J Biol Chem* 2001;276:32022–30.
- 106 Weng AP, Nam Y, Wolfe MS, Pear WS, Griffin JD, Blacklow SC, et al. Growth Suppression of Pre-T acute lymphoblastic leukemia cells by inhibition of notch signaling. *Mol Cell Biol* 2003;23:655–64.
- 107 Moeller RE, Cornejo M, Davis TN, Del Bianco C, Aster JC, Blacklow SC, et al. Direct inhibition of the NOTCH transcription factor complex. *Nature* 2009;462:182–8.
- 108 Sharff KA, Song W-X, Luo X, Tang N, Luo J, Chen J, et al. Hey1 basic helix-loop-helix protein plays an important role in mediating BMP9-induced osteogenic differentiation of mesenchymal progenitor cells. *J Biol Chem* 2009;284:649–59.
- 109 Fu Y, Chang A, Chang L, Niessen K, Eapen S, Setiadi A, et al. Differential regulation of transforming growth factor β signaling pathways by Notch in human endothelial cells. *J Biol Chem* 2009;284:19452–62.
- 110 Itoh F, Itoh S, Goumans MJ, Valdimarsdottir G, Iso T, Dotto GP, et al. Synergy and antagonism between Notch and BMP receptor signaling pathways in endothelial cells. *EMBO J* 2004;23:541–51.
- 111 Xu J, Lamouille S, Derynck R. TGF-beta-induced epithelial to mesenchymal transition. *Cell Res* 2009;19:156–72.
- 112 Kerr G, Sheldon H, Chaikwad A, Alfano I, Delft FV, Bullock AN, et al. A small molecule targeting ALK1 prevents Notch cooperativity and inhibits functional angiogenesis. *Angiogenesis* 2015;18:209–17.
- 113 Bendell JC, Gordon MS, Hurwitz HI, Jones SF, Mendelson DS, Blobe GC, et al. Safety, pharmacokinetics, pharmacodynamics, and antitumor activity of dalantercept, an activin receptor-like kinase-1 ligand trap, in patients with advanced cancer. *Clin Cancer Res* 2014;20:480–9.
- 114 Simonelli M, Zucali P, Santoro A, Thomas MB, Braud FGD, Borghaei H, et al. Phase I study of PF-03446962, a fully human monoclonal antibody against activin receptor-like kinase 1 in patients with hepatocellular carcinoma. *Ann Oncol* 2016;27:1782–7.
- 115 Necchi A, Giannatempo P, Mariani L, Farè E, Raggi D, Pennati M, et al. PF-03446962, a fully-human monoclonal antibody against transforming growth-factor β (TGF β) receptor ALK1, in pre-treated patients with urothelial cancer: An open label, single-group, phase 2 trial. *Invest New Drugs* 2014;32:555–60.
- 116 Toshihiko D, Lee KH, Kim TM, Ohtsu A, Kim TY, Ikeda M, et al. A phase I study of the human anti-activin receptor-like kinase 1 antibody PF-03446962 in Asian patients with advanced solid tumors. *Cancer Med* 2016;5:1454–63.
- 117 Goff LW, Cohen RB, Berlin J, De Braud FG, Lyshchik A, Noverasco C, et al. A phase I study of the anti-activin receptor-like kinase 1 (ALK-1) monoclonal antibody PF-03446962 in patients with advanced solid tumors. *Clin Cancer Res* 2016;22:2146–54.
- 118 Wang X, Solban N, Khanna P, Callea M, Song J, Alsop DC, et al. Inhibition of ALK1 signaling with dalantercept combined with VEGFR TKI leads to tumor stasis in renal cell carcinoma. *Oncotarget* 2016;7:41857–69.
- 119 Akhurst RJ, Hata A. Targeting the TGF β signalling pathway in disease. *Nat Rev Drug Discov* 2012;11:790–811.
- 120 Serova M, Tijeras-Raballand A, Santos CD, Albuquerque M, Paradis V, Neuzillet C, et al. Effects of TGF-beta signalling inhibition with galunisertib (LY2157299) in hepatocellular carcinoma models and in vivo whole tumor tissue samples from patients. *Oncotarget* 2015;6:348–54.
- 121 Rodon J, Carducci MA, Azaro A, Calvo E, Seoane J, Braña I, et al. First-in-human dose study of the novel transforming growth factor- β receptor I kinase inhibitor LY2157299 monohydrate in patients with advanced cancer and glioma. *Clin Cancer Res* 2015;21:553–60.
- 122 Fujiwara Y, Nokihara H, Yamada Y, Yamamoto N, Sunami K, Utsumi H, et al. Phase I study of galunisertib, a TGF-beta receptor I kinase inhibitor, in Japanese patients with advanced solid tumors. *Cancer Chemother Pharmacol* 2015;76:1–10.
- 123 David W, Cornelia W, Manfred G. Hey bHLH transcription factors. *Curr Topics Dev Biol* 2014;110:285–315.
- 124 Fischer A, Gessler M. Delta-Notch-and then? Protein interactions and proposed modes of repression by Hes and Hey bHLH factors. *Nucleic Acids Res* 2007;35:4583–96.
- 125 Ball DW. Achaete-scute homolog-1 and Notch in lung neuroendocrine development and cancer. *Cancer Lett* 2004;204:159–69.
- 126 Axelson H. The Notch signaling cascade in neuroblastoma: Role of the basic helix-loop-helix proteins HASH-1 and HES-1. *Cancer Lett* 2004;204:171–8.
- 127 Takata T, Ishikawa F. Human Sir2-related protein SIRT1 associates with the bHLH repressors HES1 and HEY2 and is involved in HES1- and HEY2-mediated transcriptional repression. *Biochem Biophys Res Commun* 2003;301:250–7.
- 128 Gould F, Harrison SM, Hewitt EW, Whitehouse A. Kaposi's sarcoma-associated herpesvirus RTA promotes degradation of the Hey1 repressor protein through the ubiquitin proteasome pathway. *J Virol* 2009;83:6727–38.
- 129 Lavery DN, Villarronga MA, Walker MM, Patel A, Belandia B, Bevan CL. Repression of androgen receptor activity by HEYL, a third member of the Hair/Enhancer-of-split-related family of Notch effectors. *J Biol Chem* 2011;286:17796–808.
- 130 Weber D, Heisig J, Kneitz S, Wolf E, Eilers M, Gessler M. Mechanisms of epigenetic and cell-type specific regulation of Hey target genes in ES cells and cardiomyocytes. *J Mol Cell Cardiol* 2015;79:79–88.
- 131 West AC, Johnstone RW. New and emerging HDAC inhibitors for cancer treatment. *J Clin Invest* 2014;124:30–9.
- 132 Falkenberg KJ, Johnstone RW. Histone deacetylases and their inhibitors in cancer, neurological diseases and immune disorders. *Nat Rev Drug Discov* 2014;13:673–91.
- 133 Dovey OM, Foster CT, Conte N, Edwards SA, Edwards JM, Singh R, et al. Histone deacetylase 1 and 2 are essential for normal T-cell development and genomic stability in mice. *Blood* 2013;121:1335–44.
- 134 Bhaskara S, Knutson SG, Chandrasekharan MB, Wilson AJ, Zheng S, Yenamandra A, et al. Hdac3 is essential for the maintenance of chromatin structure and genome stability. *Cancer Cell* 2010;18:436–47.
- 135 Suresh P, Devaraj V, Srinivas NR, Mullangi R. Review of bioanalytical assays for the quantitation of various HDAC inhibitors such as vorinostat, belinostat, panobinostat, romidepsin and chidamine. *Biomed Chromatogr* 2017;31.
- 136 Batlevi CL, Kasamon Y, Bociek RG, Lee P, Gore L, Copeland A, et al. ENGAGE-501: Phase 2 study of entinostat (SNDX-275) in relapsed and refractory Hodgkin lymphoma. *Haematologica* 2016;101:968–75.
- 137 Ruiz R, Raez LE, Rolfó C. Entinostat (SNDX-275) for the treatment of non-small cell lung cancer. *Expert Opin Investig Drugs* 2015;24:1101–9.
- 138 Knipstein J, Gore L. Entinostat for treatment of solid tumors and hematologic malignancies. *Expert Opin Investig Drugs* 2011;20:1455–67.
- 139 Yardley DA, Ismail-Khan RR, Melichar B, Lichinitser M, Munster PN, Klein PM, et al. Randomized phase II, double-blind, placebo-controlled study of exemestane with or without entinostat in postmenopausal women with locally recurrent or metastatic estrogen receptor-positive breast cancer progressing on treatment with a nonsteroidal aromatase inhibitor. *J Clin Oncol* 2013;31:2128–35.
- 140 Galli M, Salmoiraghi S, Golay J, Gozzini A, Crippa C, Pescosta N, et al. A phase II multiple dose clinical trial of histone deacetylase inhibitor ITF2357 in patients with relapsed or progressive multiple myeloma. *Ann Hematol* 2010;89:185–90.
- 141 Venugopal B, Baird R, Kristeleit RS, Plummer R, Cowan R, Stewart A, et al. A phase I study of quisinostat (JNJ-26481585), an oral hydroxamate histone deacetylase inhibitor with evidence of target modulation and antitumor activity, in patients with advanced solid tumors. *Clin Cancer Res* 2013;19:4262–72.
- 142 Banerji U, van Doorn L, Papadatos-Pastos D, Kristeleit R, Debnam P, Tall M, et al. A phase I pharmacokinetic and pharmacodynamic study of CHR-3996, an oral class I selective histone deacetylase inhibitor in refractory solid tumors. *Clin Cancer Res* 2012;18:2687–94.
- 143 Ohnuki H, Jiang K, Wang D, Salvucci O, Kwak H, Sánchez-Martín D, et al. Tumor-infiltrating myeloid cells activate Dll4/Notch/TGF- β signaling to drive malignant progression. *Cancer Res* 2014;74:2038–49.
- 144 Mendelson J, Song S, Li Y, Maru DM, Mishra B, Davila M, et al. Dysfunctional transforming growth factor- β signaling with constitutively active Notch signaling in Barrett's esophageal adenocarcinoma. *Cancer* 2012;117:3691–702.
- 145 Sjölund J, Boström AK, Lindgren D, Manna S, Moustakas A, Ljungberg B, et al. The notch and TGF- β signaling pathways contribute to the aggressiveness of clear cell renal cell carcinoma. *PLoS One* 2011;6:e23057.

- 146 Zhang J, Wang Y, Li D, Jing S. Notch and TGF- β /Smad3 pathways are involved in the interaction between cancer cells and cancer-associated fibroblasts in papillary thyroid carcinoma. *Tumor Biol* 2014;35:379–85.
- 147 Manigat L, Purow B. DDEL-16 synergistic combination of an HDAC inhibitor (HDACi) and a Notch inhibitor versus glioblastoma and melanoma cells. *Neuro-Oncol* 2015;17:v76–v7.
- 148 Gilbert CA, Daou MC, Moser RP, Ross AH. Gamma-secretase inhibitors enhance temozolomide treatment of human gliomas by inhibiting neurosphere repopulation and xenograft recurrence. *Cancer Res* 2010;70:6870–9.
- 149 Fuino L, Bali P, Wittmann S, Donapaty S, Guo F, Yamaguchi H, et al. Histone deacetylase inhibitor LAQ824 down-regulates Her-2 and sensitizes human breast cancer cells to trastuzumab, taxotere, gemcitabine, and epothilone B. *Mol Cancer Ther* 2003;2:971–84.
- 150 Ren XF, Mu LP, Jiang YS, Wang L, Ma JF. LY2109761 inhibits metastasis and enhances chemosensitivity in osteosarcoma MG-63 cells. *Eur Rev Med Pharmacol Sci* 2015;19:1182–90.
- 151 Loconte NK, Razak ARA, Ivy P, Tevaarwerk A, Leverence R, Kolesar J, et al. A multicenter phase 1 study of γ -secretase inhibitor RO4929097 in combination with capecitabine in refractory solid tumors. *Invest New Drugs* 2014;33:169–76.
- 152 Richter S, Bedard PL, Chen EX, Clarke BA, Tran B, Hotte SJ, et al. A phase I study of the oral gamma secretase inhibitor R04929097 in combination with gemcitabine in patients with advanced solid tumors (PHL-078/CTEP 8575). *Invest New Drugs* 2014;32:243–9.
- 153 Shin DY, Kim SJ, Yoon DH, Yong P, Kong JH, Kim JA, et al. Results of a phase II study of vorinostat in combination with intravenous fludarabine, mitoxantrone, and dexamethasone in patients with relapsed or refractory mantle cell lymphoma: An interim analysis. *Cancer Chemother Pharmacol* 2016;77:1–9.
- 154 Thomas A, Rajan A, Szabo E, Tomita Y, Carter CA, Scepura B, et al. A Phase I/II trial of belinostat in combination with cisplatin, doxorubicin, and cyclophosphamide in thymic epithelial tumors: A clinical and translational study. *Clin Cancer Res* 2014;20:5392–402.
- 155 Matulonis U, Berlin S, Lee H, Whalen C, Obermayer E, Penson R, et al. Phase I study of combination of vorinostat, carboplatin, and gemcitabine in women with recurrent, platinum-sensitive epithelial ovarian, fallopian tube, or peritoneal cancer. *Cancer Chemother Pharmacol* 2015;76:417–23.
- 156 Akerley W, McCoy J, Hesketh PJ, Goodwin JW, Bearden JD, Atkins JN, et al. Gemcitabine and irinotecan for patients with untreated extensive stage small cell lung cancer: SWOG 0119. *J Thorac Oncol* 2007;2:526–30.
- 157 Yoo C, Ryu MH, Na YS, Ryoo BY, Lee CW, Kang YK. Vorinostat in combination with capecitabine plus cisplatin as a first-line chemotherapy for patients with metastatic or unresectable gastric cancer: phase II study and biomarker analysis. *Br J Cancer* 2016;114:1185–90.
- 158 Fischer A, Klattig J, Kneitz B, Diez H, Maier M, Holtmann B, et al. Hey basic helix-loop-helix transcription factors are repressors of GATA4 and GATA6 and restrict expression of the GATA target gene ANF in fetal hearts. *Mol Cell Biol* 2005;25:8960–70.
- 159 Sun J, Kamei CN, Layne MD, Jain MK, Liao JK, Lee ME, et al. Regulation of myogenic terminal differentiation by the hairy-related transcription factor CHF2. *J Biol Chem* 2001;276:18591–6.
- 160 Arai MA, Masada A, Ohtsuka T, Kageyama R, Ishibashi M. The first Hes1 dimer inhibitors from natural products. *Bioorg Med Chem Lett* 2009;19:5778–81.
- 161 Müller P, Kietz S, Gustafsson J, Ström A. The Anti-estrogenic effect of all-trans-retinoic acid on the breast cancer cell line MCF-7 is dependent on HES-1 expression. *J Biol Chem* 2002;277:28376–9.
- 162 Danza G, Di Serio C, Rosati F, Lonetto G, Sturli N, Kacer D, et al. Notch signaling modulates hypoxia-induced neuroendocrine differentiation of human prostate cancer cells. *Mol Cancer Res* 2012;10:230–8.
- 163 Skinner MK, Rawls A, Wilsonrawls J, Roalson EH. Basic helix-loop-helix transcription factor gene family phylogenetics and nomenclature. *Differentiation* 2010;80:1–8.
- 164 Tsigelny IF, Kouznetsova VL, Pingle SC, Kesari S. bHLH Transcription factors inhibitors for cancer therapy: General features for in silico drug design. *Curr Med Chem* 2014;21:3227–43.
- 165 Yin X, Giap C, Lazo JS, Prochownik EV. Low molecular weight inhibitors of Myc–Max interaction and function. *Oncogene* 2003;22:6151–9.
- 166 Kiessling A, Sperl B, Hollis A, Eick D, Berg T. Selective inhibition of c-Myc/Max dimerization and DNA binding by small molecules. *Chem Biol* 2006;13:745–51.
- 167 Hammoudeh DI, Follis AV, Prochownik EV, Metallo SJ. Multiple independent binding sites for small-molecule inhibitors on the oncoprotein c-Myc. *J Am Chem Soc* 2009;131:7390–401.
- 168 Scheuermann TH, Li Q, Ma HW, Key J, Zhang L, Chen R, et al. Allosteric inhibition of hypoxia inducible factor-2 with small molecules. *Nat Chem Biol* 2013;9:271–6.
- 169 Lee KA, Zhang H, Qian DZ, Rey S, Liu JO, Semenza GL. Acriflavine inhibits HIF-1 dimerization, tumor growth, and vascularization. *Proc Natl Acad Sci U S A* 2009;106:17910–5.
- 170 Ghosh I, Chmielewski J. A β -sheet peptide inhibitor of E47 dimerization and DNA binding. *Chem Biol* 1998;5:439–45.
- 171 Chen CH, Kuo SC, Huang LJ, Hsu MH, Lung FDT. Affinity of synthetic peptide fragments of MyoD for Id1 protein and their biological effects in several cancer cells. *J Pept Sci* 2010;16:231–41.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

- Q1: Page: 1: AU: Per journal style, genes, alleles, loci, and oncogenes are italicized; proteins are roman. Please check throughout to see that the words are styled correctly. AACR journals have developed explicit instructions about reporting results from experiments involving the use of animal models as well as the use of approved gene and protein nomenclature at their first mention in the manuscript. Please review the instructions at <http://www.aacrjournals.org/site/InstrAuthors/ifora.xhtml#genenomen> to ensure that your article is in compliance. If your article is not in compliance, please make the appropriate changes in your proof.
- Q2: Page: 1: Author: Please verify the drug names and their dosages used in the article.
- Q3: Page: 1: Author: Please verify the edits to the article title for correctness.
- Q4: Page: 1: Author: Please verify the affiliations and their corresponding author links.
- Q5: Page: 1: Author: Please verify the corresponding authors' details.
- Q6: Page: 1: Author: Please check the sentence "Hairy and Enhancer-of-split related with YRPW motif factors (Hey1/2/L) belong. . ." for clarity.
- Q7: Page: 2: Author: Please confirm quality/labeling of all images included within this article. Thank you.
- Q8: Page: 2: Author: Please verify the layout of Tables 1 and 2 for correctness.
- Q9: Page: 5: Author/PE: Please reveal the unapproved structures of the compounds mentioned in Table 2 or cite the references that contain their structures.
- Q10: Page: 7: Author: Please check the sentence " For instance, 64% thymoma and thymic carcinoma patients show objective. . ." for clarity.
- Q11: Page: 8: AU:/PE: The conflict-of-interest disclosure statement that appears in the proof incorporates the information from forms completed and signed off on by each individual author. No factual changes can be made to disclosure information at the proof stage. However, typographical errors or misspelling of author names should be noted on the proof and will be corrected before publication. Please note if any such errors need to be corrected. Is the disclosure statement correct?
- Q12: Page: 8: Author: Please verify the headings "Acknowledgments" and "Grant Support" and their content for correctness.
- Q13: Page: 8: Author: Refs. 13, 34, 48, 68, 71, 73, 76, 87, 92, 103, 114, 117, 118, 143, and 157 have been updated as per PubMed. Please verify.
- Q14: Page: 9: Author: Please provide page range for refs. 46 and 135.
- Q15: Page: 10: Author: Ref. 92 has been updated as per PubMed. Please verify.

Q16: Page: 10: Author: Ref. 93 has been updated as per "http://cancerres.aacrjournals.org/content/73/8_Supplement/213." Please verify.

AU: Below is a summary of the name segmentation for the authors according to our records. The First Name and the Surname data will be provided to PubMed when the article is indexed for searching. Please check each name carefully and verify that the First Name and Surname are correct. If a name is not segmented correctly, please write the correct First Name and Surname on this page and return it with your proofs. If no changes are made to this list, we will assume that the names are segmented correctly, and the names will be indexed as is by PubMed and other indexing services.

First Name	Surname
Zihao	Liu
Andrew J.	Sanders
Gehao	Liang
Erwei	Song
Wen G.	Jiang
Chang	Gong